



Ordering physician:
Carrie Jones

DOB:1974-11-13
Gender: Male

Collection Times:
2016-01-12 08:20PM
2016-01-12 10:56PM
2016-01-13 05:30AM
2016-01-13 07:58AM

Hormone Testing Summary

All units are given in ng/mg creatinine

low limit * your result * high limit

How to read the graphical representation of results

Sex Hormones

14.3

Total Estrogen
(Sum of 8 Estrogen Metabolites)

61.0

Testosterone

See Pages 2 & 3 for a thorough breakdown of sex hormone metabolites

	Age	Range
Testosterone	20-40	40-100
	40-60	30-60
	>60	10-40
Total DHEA Production	20-40	2500-5500
	40-60	1700-3500
	>60	1000-2500

Adrenal Hormones

See pages 4 and 5 for a more complete breakdown of adrenal hormones

Daily Free Cortisol Pattern

Cortisol (ng/mg)

High Range Limit

Low Range Limit

Patient Values

Waking (A) Morning (B) Afternoon (C) Night (D)

2239.0

Total DHEA Production
(DHEAS + Etiocholanolone + Androsterone)

115.0

24hr Free Cortisol
(A+B+C+D)

cortisol metabolism

5497.0

Metabolized Cortisol (THF+THE)
(Total Cortisol Production)

Free cortisol best reflects tissue levels. Metabolized cortisol best reflects total cortisol production.

Patient Reported Hormone Therapies: ROA 1=oral, 2=sublingual, 3=transdermal cream, 4=transdermal gel, 5=vaginal/labial, 6=rectal mucosa, 7=patch, 8=pellet, 9=injection, 10=other

Hormone	Brand	ROA (1-10)	Dose (mg)	Date Last Used	Times per Day	Length of use
The patient did not report any relevant hormone therapies						

TO THE LEFT YOU CAN SEE A SCREENSHOT OF THE THERAPY SECTION OF THE PATIENT REQUISITION.

There is a series of videos in our video library at dutchtest.com that you may find useful in understanding the report. The following videos (which can also be found on the website under the listed names) may be particularly helpful in aiding your understanding:

- [Analytical Report Overview \(quick overview\)](#)
- [Estrogen tutorial](#) ; [Androgen tutorial](#)
- [Cortisol tutorial](#)



Dried Urine Test for Comprehensive Hormones

Sex Hormones and Metabolites

Accession # 00219441

Damien Blenkinsopp
30 Moorland Road
Hemel Hempstead,




Ordering physician:
Carrie Jones

DOB:1974-11-13
Gender: Male

Collection Times:
2016-01-12 08:20PM
2016-01-12 10:56PM
2016-01-13 05:30AM
2016-01-13 07:58AM

Category	Test	Result	Units	Normal Range
Progesterone Metabolism				
	b-Pregnanediol	Within range	180.0	ng/mg 40 - 300
	a-Pregnanediol	Within range	55.0	ng/mg 15 - 110
Androgen Metabolism				
	DHEAS	Low end of range	81.0	ng/mg 60 - 2000
	Androsterone	Within range	1493.0	ng/mg 640 - 3000
	Etiocholanolone	Low end of range	665.0	ng/mg 460 - 1700
	Testosterone	Within range	61.0	ng/mg 25 - 100
	5a-DHT	Low end of range	10.1	ng/mg 9 - 16.7
	5a-Androstanediol	Within range	111.6	ng/mg 49 - 147
	5b-Androstanediol	Low end of range	43.9	ng/mg 30 - 147
	Epi-Testosterone	Within range	76.6	ng/mg 25 - 100
Estrogen Metabolites				
	Estrone(E1)	Within range	6.3	ng/mg 4 - 12
	Estradiol(E2)	Within range	0.9	ng/mg 0.5 - 1.6
	Estriol(E3)	Low end of range	2.6	ng/mg 2 - 6
	2-OH-E1	Above range	3.6	ng/mg 0 - 3
	4-OH-E1	Within range	0.3	ng/mg 0 - 0.5
	16-OH-E1	Within range	0.6	ng/mg 0 - 0.8
	2-Methoxy-E1	Within range	0.8	ng/mg 0 - 2
	2-OH-E2	Within range	0.26	ng/mg 0 - 0.5

HOW TO READ YOUR RESULTS: Hormones are presented on this page graphically in the order the body metabolizes them. Arrows represent conversion from one hormone to another. The stars represent the low and high limits of the reference ranges (see example, right). The number in the middle is your result.





Dried Urine Test for Comprehensive Hormones

Advanced Adrenal Assessment

Accession # 00219441

Damien Blenkinsopp
30 Moorland Road
Hemel Hempstead,





Ordering physician:
Carrie Jones

DOB:1974-11-13
Gender: Male


Collection Times:
2016-01-12 08:20PM
2016-01-12 10:56PM
2016-01-13 05:30AM
2016-01-13 07:58AM

Category	Test		Result	Units	Normal Range
Creatinine					
	Creatinine A (Waking)	Within range	1.42	mg/ml	0.3 - 3
	Creatinine B (Morning)	Within range	1.66	mg/ml	0.3 - 3
	Creatinine C (Afternoon)	Within range	1.32	mg/ml	0.3 - 3
	Creatinine D (Night)	Within range	1.19	mg/ml	0.3 - 3
Daily Free Cortisol and Cortisone					
	Cortisol A (Waking)	Below range	9.6	ng/mg	17 - 50
	Cortisol B (Morning)	Within range	92.5	ng/mg	43 - 130
	Cortisol C (Afternoon)	Below range	9.2	ng/mg	17 - 34
	Cortisol D (Night)	Low end of range	3.4	ng/mg	0 - 17
	Cortisone A (Waking)	Below range	35.0	ng/mg	46 - 90
	Cortisone B (Morning)	High end of range	189.7	ng/mg	93 - 210
	Cortisone C (Afternoon)	Below range	39.5	ng/mg	40 - 90
	Cortisone D (Night)	Within range	9.8	ng/mg	0 - 45
	24hr Free Cortisol	Low end of range	115.0	ug	100 - 225
	24hr Free Cortisone	Low end of range	274.0	ug	250 - 400
Cortisol Metabolites and DHEAS					
	a-Tetrahydrocortisol (a-THF)	Within range	365.0	ng/mg	220 - 720
	b-Tetrahydrocortisol (b-THF)	Above range	2676.0	ng/mg	1330 - 2330
	b-Tetrahydrocortisone (b-THE)	Low end of range	2456.0	ng/mg	2100 - 4000
	Metabolized Cortisol (THF+THE)	Within range	5497.0	ng/mg	3850 - 7000
	DHEAS	Low end of range	81.0	ng/mg	60 - 2000
Melatonin (*measured as 6-OH-Melatonin-Sulfate)					
	Melatonin* (Waking)	Below range	9.3	ng/mg	10 - 50



Total DHEA Production

Age	Range
20-40	2500-5500
40-60	1700-3500
>60	1000-2500



Adrenal Gland


DHEA

A patient's catabolic vs anabolic balance can be estimated by observing relative DHEA (anabolic) vs cortisol (catabolic) production


Cortisol

Cortisol Metabolism






Circulating Free Cortisol



Total DHEA Production (DHEAS + Etiocholanolone + Androsterone)



Metabolized Cortisol (THF+THE) (Total Cortisol Production)



Provider Notes

Thank you for testing with us! If this is your first report, you are encouraged to watch our educational videos on how to read the report. There are hyperlinks to these videos on the first page of a DUTCH Complete or in these comments (below). The videos can also be seen by going to www.DutchTest.com and visiting the video library. Comments in the report that are specific to the patient ARE IN ALL CAPS. The other information is general information that we hope you will find useful in understanding the patient's results. Reference ranges updated 7/23/2015.

THE PATIENT REPORTED SIGNIFICANT FATIGUE IN THE AFTERNOON/EVENING, BUT NOT IN THE MORNING.

Androgen Metabolism: Testosterone is made in the testis and the adrenal glands however in men the adrenal production is a somewhat insignificant fraction. a-DHT (a-dihydrotestosterone) is the most potent androgen (3X more than testosterone), but it is primarily made within the liver and target cells (it is a paracrine hormone) and not by the gonads. a-DHT is subsequently deactivated to a-androstenediol within target tissues and then conjugated (glucuronidation) for excretion. As such, a-androstenediol may better represent a-DHT even though its metabolic precursor is more biologically active and well known however we do report both on the test. Only a fraction of a-DHT formed actually enters circulation as a-DHT (Toscano, 1987). The corresponding beta metabolites (for example b-DHT) are not androgenic.

Androgens help to promote proper sexual desire and function, and generally contribute to attributes that are typically more pronounced in males than females (general and sexual aggression, muscle mass, increased facial/body hair, reduction of fat deposition, etc). As males age, they make less testosterone. Androgens are turned into estrogens (for men and women) so assessing metabolism also includes looking at estrogen levels.

5a-Reductase Activity: The competing enzymes 5a and 5b-reductase act on the androgens androstenedione (creating androsterone and etiocholanolone located under the progesterone picture) and testosterone (creating a-DHT and b-DHT). They also metabolize progesterone, and cortisol. The alpha metabolites of androstenedione and testosterone are far more androgenic than their beta counterparts. Consequently, increased 5a-reductase activity may be accompanied by clinical signs of androgenicity (excess facial hair growth, scalp hair loss, acne, irritability, oily skin, prostate issues in men...etc). If the patient heavily favors the 5a pathway and there are concerns of excess androgenicity (or prostate cancer risk), this may be worth addressing.

Estrogen Metabolism: While usually considered "female" hormones, estrogens are present in males also. In men elevated estrogen levels have been associated with weight gain, breast development, mood swings, cardiovascular and prostate problems. The primary concern for male patients is if they are making too much. Estrogens are produced from androstenedione and testosterone via a process called aromatization.

Most estrogen is metabolized down into the healthier 2-OH pathway. These 2-OH metabolites are considered protective in women (from breast cancer) but this is also true of men with respect to prostate cancer. It is ideal if the 2-OH pathway (from the 2nd pie chart) is heavily favored over the more potent 16-OH estrogens or the carcinogenic 4-OH estrogens. It is our position that the ratio of 2:16 OHE1 is not as relevant as has been thought historically (Obi, 2011). Providers may still wish to use this index and it can be calculated by simply dividing the two numbers. A male reference range for the ratio with our methodology is 2.6-6.6.

The methylation activity gauge shows how effectively the body makes methoxy estrogens from hydroxyestrogens. Normal or high methylation activity is important in protecting the body from the harmful 4-OH estrogens made in phase 1 detoxification.

Progesterone levels are of marginal value in men, although deficiency can be associated with some clinical conditions such as depression, fatigue, and low libido.

DUTCH Adrenal: The HPA-Axis refers to the communication and interaction between the hypothalamus (H) and pituitary (P) in the brain down to the adrenal glands (A) that sit on top of your kidneys. When a physical or psychological stressor occurs, the hypothalamus tells the pituitary to make ACTH, a hormone. ACTH stimulates the adrenal glands to make the stress hormone, cortisol and to a lesser extent DHEA and DHEA-S. Normally, the HPA-axis production follows a daily pattern in which cortisol rises rather rapidly in the first 10-30 minutes after waking in order to help with energy, then gradually decreases throughout the day so that it is low at night for sleep. The cycle starts over the next morning. Abnormally high activity occurs in Cushing's Disease where the HPA-axis is hyper-stimulated causing cortisol to be elevated all day. The opposite is known as Addison's Disease, where cortisol is abnormally low because it is not made appropriately in response to ACTH's stimulation. These two conditions are somewhat rare. Examples of more common conditions related to less severely abnormal cortisol levels include fatigue, depression, insomnia, fibromyalgia, anxiety, inflammation and more.

Only a fraction of cortisol is "free" and bioactive. This fraction of cortisol is very important, but levels of metabolized cortisol best represents overall production of cortisol therefore both should be taken into account to correctly assess adrenal function.

The Daily Free Cortisol Pattern: In healthy adrenal function, cortisol levels are expected to rise in the early morning and fall throughout the day, reaching the lowest point around 1am and peaking 30-60min after waking. The waking sample represents the total of free cortisol throughout the sleeping period. Cortisone is the inactive form of cortisol. Its pattern is of secondary importance, but at times can give additional clarity and is provided on the adrenal page. Typical urine testing (24-hour collection) averages the daily production of cortisol. This approach is not able to properly characterize individuals whose

cortisol patterns do not fit the typical rise then fall pattern through the day. Dysfunctional diurnal patterns have been associated with health-related problems such as fatigue and insomnia.

The daily total of free cortisol is approximated by summing the four measurements. This calculated value correlates to a 24-hour free cortisol value. It is helpful to compare the relative level of 24-hr free cortisol with metabolized cortisol to understand HPA-axis activity. The total of free cortisol for the day only represents about 1-3% of the total production. The total of the metabolites is a better marker for overall cortisol production.

FREE CORTISOL LEVELS ARE ON THE LOWER SIDE OF THE REFERENCE RANGE. LEVELS OF METABOLIZED CORTISOL CONFIRM THAT OVERALL CORTISOL PRODUCTION IS REASONABLE, AND THE ACTUAL DIURNAL PATTERN OF FREE CORTISOL SHOULD BE EXAMINED TO FURTHER EXAMINE CORTISOL PRODUCTION.

The Cortisol-Cortisone Balance: Cortisol, which is the active hormone, can convert into cortisone, the inactive form. They convert back and forth in different parts of the body. We tell which one you make more of by looking at whether cortisol metabolites (aTHF, bTHF) or cortisone metabolites (bTHE) are made more (compared to what is normal) in the gauge at the bottom of the adrenal page. The deactivation of cortisol to cortisone (via enzyme 11b-HSD II) occurs predominantly in the kidneys, colon, and saliva glands. The local formation of inactive cortisone from cortisol in the kidney is strongly reflected in urine. Activation of cortisone to cortisol takes place primarily in the liver, adipose tissue, gonads, brain, and muscle. Within these same tissues (mostly the liver) the free hormones are also converted to their metabolites (cortisol to a/b-THF, cortisone to THE). Balance between the two is usually preferred, but making more cortisol than cortisone is sometimes good to help give you enough cortisol if your levels are low however a preference for the active cortisol is enhanced by central adiposity, hypothyroidism, inflammation, and supplements such as licorice root extract. Cortisone formation is enhanced by growth hormone, estrogen, coffee and hyperthyroidism.

Reading the Report: The first page of the Dutch Complete lab report is a summary page while the second page of the Dutch Complete lab report and first page of the Dutch sex hormone and Dutch adrenal test are a classic lab report showing each result and the respective range of each hormone. Reference ranges shown are those of young healthy individuals. The graphical representation of results on the page that follows allows the viewing of hormone results within the biochemical flowchart to more easily see the relative level of each hormone. The gauge format shows the patient result (represented by the "needle" of the gauge) and the area between the stars represents the reference range. Reference ranges are typically set at the 20th to the 80th percentile of young, healthy individuals (DHEAS for example). This means that a result at the low end of a range is lower than 80 percent of young, healthy individuals. Likewise a result at the high end of a range is higher than 80 percent of the population. Some reference ranges are set more widely. For example, slightly elevated progesterone is not generally considered problematic, so its metabolites have reference ranges that extend further (90th percentile instead of 80th).

The "fan" style gauges are used for indexes/ratios such as on 5a-reductase activity, cortisol/cortisone, and estrogen methylation. Because these values are all based on ratios there are no values or units, but they give a general idea of a particular relationship and can tell you how 'turned up' or 'turned down' a particular process is. The middle of the gauge represents an average value, while the lines towards the edge represent results lower or higher than most (80%) of the population. Being outside of any range is not always considered unfavorable. For example, on the estrogen methylation gauge an elevated level means someone methylates estrogens very effectively which may have positive consequences.

What is actually measured in urine? In blood, most hormones are bound to binding proteins. A small fraction of the total hormone levels are "free" and unbound such that they are active hormones. These free hormones are not found readily in urine except for cortisol and cortisone (because they are much more water soluble than, for example, testosterone). As such, free cortisol and cortisone can be measured in urine and it is this measurement that nearly all urinary cortisol research is based upon. In the DUTCH Adrenal Profile the diurnal patterns of free cortisol and cortisone are measured by LC-MS/MS.

All other hormones measured (cortisol metabolites, DHEA, and all sex hormones) are excreted in urine predominately after the addition of a glucuronide or sulfate group (to increase water solubility for excretion). As an example, Tajic (Natural Sciences, 1968 publication) found that of the testosterone found in urine, 57-80% was testosterone-glucuronide, 14-42% was testosterone-sulfate, and negligible amounts (<1% for most) was free testosterone. The most likely source of free sex hormones in urine is from contamination from hormonal supplements. To eliminate this potential, we remove free hormones from conjugates. The glucuronides and sulfates are then broken off of the parent hormones, and the measurement is made. These measurements reflect well the bioavailable amount of hormone in most cases as it is only the free, nonprotein-bound fraction in blood/tissue that is available for phase II metabolism (glucuronidation and sulfation) and subsequent urine excretion.

Disclaimer: the filter paper used for sample collection is designed for blood collection, so it is technically considered "research only" for urine collection. Its proper use for urine collection has been thoroughly validated.